

Bioefficacy of Pyroxsulam (XDE-742) for Weed Control in Wheat (*Triticum aestivum* L.)

V. Pratap Singh, V. C. Dhyani, S. P. Singh, Abnish Kumar, M. K. Singh and Neeta Tripathi

Department of Agronomy

G. B. Pant University of Agriculture & Technology, Pantnagar-263 145 (Uttarakhand), India

The acute problem of both grassy and broad leaf weeds is becoming very common in wheat growing areas of north-western zone of India, which often results in huge yield losses and makes the weed control more complex (Singh *et al.*, 2002). Several herbicides have been used from time to time to control weeds in wheat. Continuous use of the same herbicide or herbicides having the same mode of action may result in shift in weed flora, development of resistance in weeds (Moss and Rubin, 1993) as well as build up of residue in soil. Therefore, a new herbicide molecule pyroxsulam (XDE-742) was used in the present experiment during rabi 2005-06. Pyroxsulam is a broad spectrum herbicide which controls grassy as well as broad leaf weeds in wheat.

The present study was carried out at Crop Research Centre of G. B. Pant University of Agriculture & Technology, Pantnagar in a randomized block design with three replications. Fifteen treatments with different doses and concentrations of Pyroxsulam (12, 15, 18 and 30 g/ha of 3.0 and 3.6% O. D. both) alongwith 2, 4-D ethyl ester 190 g/ha and aminopyralid 7.5 g/ha were taken in the experimental plot. Treatments having sulfosulfuron 25 g/ha, clodinafop-propargyl 60 g/ha and isoproturon 1000 g/ha, weed free and weedy plot were also included as standard check. The soil of the experimental field was silty clay loam, high in organic carbon (0.76%), medium in available phosphorus (19 kg/ha) and potassium (225 kg/ha) with pH 7.1. The recommended doses of fertilizer i. e. 120 : 60 : 40 kg NPK/ha were applied in the experimental plot. Wheat variety PBW 343 was sown on 12 December 2005. Half of nitrogen and full doses of P and K were applied as basal dressing before sowing. The remaining half dose of nitrogen was top-dressed into two equal splits at tillering and heading stage of wheat. Herbicides were applied 30 days after sowing through knapsack sprayer using 600 l/ha water.

Weed counts were recorded at 30 days after herbicide application by taking the observation with the help of quadrat of 0.25 m² from each plot and weed dry

weight was recorded by keeping the sample in oven at 70 ± 1°C for 48 h.

The experimental plot was mainly infested with *Phalaris minor*, *Melilotus indica*, *Coronopus didymus*, *Lathyrus aphaca* and *Chenopodium album* which account for 26, 22, 20, 10 and 7%, respectively, in weedy plot at 30 days after sowing (Table 1). As the dose increased from 12 to 30 g, significant reduction in weed density of *P. minor* was observed with pyroxsulam. Application of pyroxsulam at 12 and 15 g recorded significantly lower weed density of *P. minor* at their lower concentration (3.0% O. D.) as compared to their higher concentration (3.6% O. D.). However, pyroxsulam at their higher doses (18 and 30 g) recorded similar density of *P. minor* at both the concentrations at 30 days after herbicide application. Clodinafop was found effective against *P. minor* but not against other weeds. The density of *L. aphaca* and *M. indica* reduced due to increase in doses of pyroxsulam. The efficacy of pyroxsulam increased against these weeds when applied with aminopyralid at 7.5 g/ha. Isoproturon at 1.0 kg/ha also provided good control of *L. aphaca* and *M. indica* but its efficacy was poor on *P. minor*, *C. didymus* and *C. album*. Pyroxsulam was found effective to control the *C. didymus*, *C. album* and *M. denticulata* which was evident from their zero weed density by pyroxsulam over the standard herbicide check viz., sulfosulfuron, clodinafop and isoproturon. With increase in the dose of herbicide, there was decrease in the total weed dry weight at 30 days after herbicide application at both the concentrations. Highest weed control efficiency was observed in weed free situation. It was followed by pyroxsulam supplemented with aminopyralid (87.5) which recorded similar weed control efficiency as that of pyroxsulam at 30 g/ha at both the concentrations.

Application of pyroxsulam (12 to 30 g) increased the grain yield in both the concentrations (3.0 and 3.6% O.D.); however, the differences among the doses were non-significant (Table 2). Pyroxsulam without intron alongwith 2, 4-D gave similar grain yield (48.17 q/ha) as with weed free (48.18 q/ha) situation. Lower yield at

Table 1. Effect of different treatments on weed species (No./m²) and total weed dry weight (g/m²) at 30 days after transplanting

Treatments	Dose (g a. i/ha)	<i>P. minor</i>	<i>Cornopus didymus</i>	<i>Laythyrus aphaca</i>	<i>Chenopodium album</i>	<i>Melilotus indica</i>	<i>Medicago sativa</i>	<i>Vicia sativa</i>	<i>Polygonum plebejum</i>	Total weed dry weight (g/m ²)	Weed control efficiency (WCE)
XDE 742 3% OD+Intron	12+300	(36.7)	(0.0)	(14.0)	(0.0)	(10.0)	(0.0)	(3.3)	(4.0)	12.1	75.2
		3.6	0.0	2.6	0.0	2.4	0.0	1.4	1.6		
XDE 742 3% OD+Intron	15+300	(27.3)	(0.0)	(11.3)	(0.0)	(5.3)	(0.0)	(1.3)	(0.7)	8.5	82.5
		3.3	0.0	2.5	0.0	1.7	0.0	0.7	0.4		
XDE 742 3% OD+Intron	18+300	(24.0)	(0.0)	(8.7)	(0.0)	(4.0)	(0.0)	(0.0)	(0.7)	7.7	84.2
		3.2	0.0	2.3	0.0	1.5	0.0	0.0	0.4		
XDE 742 3% OD+Intron	30+300	(20.7)	(0.0)	(5.3)	(0.0)	(2.0)	(0.0)	(0.0)	(0.0)	6.2	87.3
		3.0	0.0	1.8	0.0	0.9	0.0	0.0	0.0		
XDE 742 3.6% OD+Intron	12+300	(59.3)	(0.0)	(13.3)	(0.0)	(14.7)	(0.0)	(4.7)	(2.0)	21.6	55.7
		4.1	0.0	2.6	0.0	2.7	0.0	1.6	0.9		
XDE 742 3.6% OD+Intron	15+300	(38.7)	(0.0)	(12.0)	(0.0)	(11.3)	(0.0)	(4.0)	(2.0)	16.5	66.2
		3.7	0.0	2.5	0.0	2.4	0.0	1.5	0.9		
XDE 742 3.6% OD+Intron	18+300	(28.7)	(0.0)	(10.7)	(0.0)	(4.0)	(0.0)	(3.3)	(0.0)	9.2	81.1
		3.4	0.0	2.4	0.0	1.3	0.0	1.4	0.0		
XDE 742 3.6% OD+Intron	30+300	(20.0)	(0.0)	(8.7)	(0.0)	(2.7)	(0.0)	(2.7)	(0.0)	6.5	86.7
		3.0	0.0	2.2	0.0	0.7	0.0	1.3	0.0		
XDE 742 3% OD+2, 4-DEE 38 EC	15+190	(24.0)	(0.0)	(14.0)	(0.0)	(2.0)	(0.0)	(0.7)	(1.3)	7.3	85.0
		3.2	0.0	2.0	0.0	0.9	0.0	0.4	0.5		
XDE 742 3% OD+Aminopyralid+Intron	15+7.5+300	(26.7)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	6.1	87.5
		3.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
Sulfosulfuron 75 WDG+Safener	25	(23.3)	(19.3)	(9.3)	(2.0)	(8.7)	(0.0)	(4.7)	(0.7)	16.9	65.4
		3.1	2.9	1.7	0.9	2.2	0.0	1.7	0.4		
Clodinafop-propargyl 15 WP	60	(0.0)	(30.7)	(22.7)	(24.7)	(38.7)	(8.7)	(9.3)	(11.3)	25.3	48.2
		0.0	3.4	3.1	3.2	3.6	2.3	2.3	2.4		
Isoproturon	1000	(26.7)	(15.3)	(4.0)	(13.3)	(5.3)	(7.3)	(2.7)	(10.7)	18.7	61.7
		3.3	2.8	1.3	2.6	1.4	2.1	1.0	2.4		
Weed free	-	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	0.0	100.0
		0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
Weedy	-	(62.67)	(48.7)	(17.3)	(24.0)	(52.7)	(10.0)	(14.7)	(12.0)	48.8	
		4.1	3.8	2.8	3.1	3.8	2.4	2.5	2.5		
LSD (P=0.05)	-	0.4	0.2	1.2	0.4	1.0	0.1	0.8	0.8	5.12	

Values in parentheses are original and transformed to log (x+1) for analysis.

Table 2. Effect of different treatments on yield and yield attributing characteristics of wheat

Treatments	Dose (g/ha)	Panicles ² (No./m ²)	No. of grains/panicle	1000-grain weight (g)	Grain yield (q/ha)	Straw yield (q/ha)
XDE 742 3% OD+Intron	12+300	403	38.40	38.63	43.34	77.52
XDE 742 3% OD+Intron	15+300	424	41.60	40.30	44.17	81.68
XDE 742 3% OD+Intron	18+300	423	36.35	37.80	44.73	89.60
XDE 742 3% OD+Intron	30+300	463	39.15	37.35	46.26	88.98
XDE 742 3.6% OD+Intron	12+300	406	37.80	40.25	41.11	75.02
XDE 742 3.6% OD+Intron	15+300	423	34.35	42.65	42.09	76.89
XDE 742 3.6% OD+Intron	18+300	423	36.35	40.15	42.65	76.89
XDE 742 3.6% OD+Intron	30+300	438	34.73	40.10	43.06	79.19
XDE 742 3% OD+2, 4-DEE38 EC	15+190	461	37.45	41.95	49.17	87.10
XDE 742 3% OD+Aminopyralid+Intron	15+7.5+300	467	39.00	41.65	48.48	85.85
Sulfosulfuron 75 WDG+Safener	25	479	30.75	41.00	47.79	94.81
Clodinafop-propargyl 15 WP	60	407	36.50	37.60	38.75	61.89
Isoproturon	1000	413	35.00	37.55	38.61	67.52
Weed free	-	415	35.00	38.80	49.18	89.39
Weedy	-	335	36.45	34.25	34.31	56.68
LSD (P=0.05)	-	65.6	NS	NS	5.09	12.60

NS–Not Significant

higher concentration (3.6%) of pyroxsulam was mainly attributed to their higher weed dry matter at 30 days after herbicide application. The highest reduction (30%) of grain yield was recorded in weedy plot over the weed free situation. Among the yield attributes, significant differences were obtained in number of panicles/m²; however, the number of grains/panicle and 1000-grain weight had non-significant difference with application of herbicide either applied as alone or in combination. Sulfosulfuron at 25 g/ha recorded the highest number of panicles/m² (479), it was followed by pyroxsulam at 15 g/ha followed by aminopyralid (467) which recorded the higher grain (48.48 q/ha) as compared to sulfosulfuron mainly due to more number of grains per

panicle. Thus, it was concluded that pyroxsulam was found effective against most of the weeds. pyroxsulam alongwith 2, 4-D recorded the similar yield in weed free situation followed by application of pyroxsulam 3%+aminopyralid+intron.

REFERENCES

- Moss, S. R. and B. Rubin. 1993. Herbicide-resistant weeds : a worldwide perspective. *J. Agric. Sci.* **120** : 141-148.
- Singh, G., V. P. Singh and M. Singh. 2002. Bioefficacy of metsulfuron-methyl in combination with isoproturon for control of grassy and non-grassy weeds in wheat. *Ind. J. Weed Sci.* **34** : 9-12.